

CORE 3

SUMMARY NOTES

1 Functions

- A function is a rule which generates exactly ONE OUTPUT for EVERY INPUT. To be defined fully the function has
 - a RULE – tells you how to calculate the output from the input
 - a DOMAIN – the set of values which will be used as inputs

e.g. $f(x) = \sqrt{x}$ domain $x \geq 0$
 (cannot find the square root of negative values)


- ALTERNATIVE NOTATION

$f: x \mapsto x^2$ means function f such that x maps to x^2
 input x is converted to output x^2

$x \in \mathbb{R}$ x can be any real number

- There are different types of functions


MANY-ONE



$$y = x^2$$

2 different inputs give the same output

ONE-ONE



$$y = 2x + 3$$

for each output there is only one possible input

- The RANGE of a function is the complete set of all of the OUTPUTS
- An INVERSE function is denoted by f^{-1} .

ONLY ONE-ONE FUNCTIONS HAVE INVERSES

The DOMAIN of an inverse function is the RANGE of the function

e.g. The function f is defined by $f(x) = \frac{3}{2x-1}$ find $f^{-1}(x)$

Step 1 : Write the rule in terms of x and y

$$y = \frac{3}{2x-1}$$

Step 2 : Rearrange to make x the subject

$$x = \frac{3+y}{2y}$$

Step 3 : Replace the y 's with x 's

$$f^{-1}(x) = \frac{3+x}{2x}$$

- Using the same scale on the x and y axis, the graphs of a function and its inverse have **reflection symmetry** in the line $y = x$

COMPOSITE FUNCTIONS

The function **gf** is called a **composite** function and tells you to 'do f first then gf(x)

e.g. $f(x) = 2x + 3$ $g(x) = x^2 + 2$

$$gf(x) = (2x+3)^2 + 2$$

$$gf(x) = 4x^2 + 12x + 11$$

$$fg(x) = 2(x^2 + 2) + 3$$



$$fg(x) = 2x^2 + 7$$

2 The Modulus function

- $|x|$ is the 'modulus of x' or the 'absolute value'
- The modulus of a real number can be thought of as its 'distance' from 0 and it is always positive.

$$|4| = 4 \quad |-2| = 2$$

- The graph of $y = |f(x)|$ is




To sketch the graph of $y = |f(x)|$ first sketch the graph of $y = f(x)$. Take any part of the graph that is below the x-axis and reflect it in the x-axis.

SOLVING EQUATIONS

Always sketch the graph before you start to determine the number of solutions

A function is defined by $f(x) = |2x+1| - 3$
Solve the inequality $f(x) < x$



The graphs shows 2 solutions

$$(2x+1) - 3 = x \quad - (2x + 1) - 3 = x$$

$$2x - 2 = x \quad -2x - 4 = x$$

$$x = 2 \quad \text{or} \quad x = -\frac{4}{3}$$

3 Transforming Graphs

- TRANSLATION - to find the equation of a graph after a translation of $\begin{bmatrix} a \\ b \end{bmatrix}$ you replace x by $(x-a)$ and y by $(y - b)$

e.g. The graph of $y = x^2 - 1$ is translated through $\begin{bmatrix} 3 \\ -2 \end{bmatrix}$. Write down the equation of

$$y - b = f(x-a)$$


or

$$y = f(x-a) + b$$

the graph formed.

$$(y + 2) = (x-3)^2 - 1$$

$$y = x^2 - 6x + 6$$



- REFLECTING

Reflection in the x -axis, replace y with $-y$
 Reflection in the y -axis, replace x with $-x$

$y = -f(x)$

$y = f(-x)$

- STRETCHING

Stretch of factor k in the x direction replace x by $\frac{1}{k}x$

$y = f\left(\frac{1}{k}x\right)$

Stretch of factor k in the y direction replace y by $\frac{1}{k}y$

$y = kf(x)$

- COMBINING TRANSFORMATIONS

When applying 2 transformations the order does not matter if one involves replacing x and the other replacing y . If both transformations involve replacing x (or y) then the order could matter

e.g. The graph of $y = x^2$ is first translated by $\begin{bmatrix} 3 \\ 0 \end{bmatrix}$ and then reflected in the y -axis

Find the equation of the final image.

Translation $y = (x - 3)^2$

Reflection $y = (-x - 3)^2$


$y = (x + 3)^2$

4 TRIGONOMETRY INVERSE FUNCTIONS


$y = \sin^{-1} x$ $\arcsin x$ or $\text{asin } x$

domain $-1 \leq x \leq 1$



range $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$




$y = \cos^{-1} x$ arccos x or acos x
 domain $-1 \leq x \leq 1$
 range $0 \leq y \leq \pi$



$y = \tan^{-1} x$ arctan x or atan x
 domain $x \in \mathbb{R}$
 range $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$



sec x is defined as $\frac{1}{\cos x}$
 $y = \sec x$ has domain $x \in \mathbb{R}$ $x \neq \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \pm \frac{5\pi}{2}$
 and range $y \leq -1$ and $y \geq 1$



cosec c is defined as $\frac{1}{\sin x}$
 $y = \text{cosec } x$ has domain $x \in \mathbb{R}$ $x \neq 0, \pm \pi, \pm 2\pi, \pm 3\pi$
 and range $y \leq -1$ and $y \geq 1$

5 Natural Logarithms and e^x

- e is an irrational; its value is 2.718281828 correct to 9 decimal places
- Natural Logarithms** use e as a base and we write $\log_e x$ as $\ln x$

$$e^x = y \Rightarrow x = \ln y$$

e.g. Solve the equation $e^{-5x} - 3 = 0$

$$e^{-5x} = 3$$

$$-5x = \ln 3$$


$$x = -\frac{1}{5} \ln 3$$

$$= -0.2197$$

If the question asks for an **exact** answer do not change into decimals

- $e^x = y \Rightarrow x = \ln y$ so e^x and $\ln x$ are inverse functions

e^x is positive for all x so
 $\ln x$ is defined only for
 positive values of x



e.g. The function g is defined by $g(x) = 2e^{x-5} + 3$ for all real x . Find an expression for $g^{-1}(x)$ and state its domain and range.

$$\begin{aligned} y &= 2e^{x-5} + 3 \\ y - 3 &= 2e^{x-5} \\ \frac{1}{2}(y - 3) &= e^{x-5} \\ \ln\left(\frac{1}{2}(y - 3)\right) &= x - 5 \\ \ln\left(\frac{1}{2}(y - 3)\right) + 5 &= x \\ g^{-1}(x) &= \ln\left(\frac{1}{2}(x - 3)\right) + 5 \end{aligned}$$

The range of g is $g(x) > 3$ so the domain of $g^{-1}(x)$ is $x > 3$

The domain of g is all real values of x so the range $g^{-1}(x)$ is all real values

- Transformation of graphs

e.g. Describe the sequence of geometrical transformations needed to obtain the graph of $y = 2e^{-x}$ from the graph of $y = e^x$.

Reflection in the y -axis gives $y = e^{-x}$

Stretch factor of 2 in the y direction gives $y = 2e^{-x}$

6 Differentiation

Key points from C1 and C2

- The derivative of $x^n = nx^{n-1}$
- If $f'(a) > 0$, f is increasing at $x = a$. If $f'(a) < 0$, f is decreasing at $x = a$
- The points where $f'(a) = 0$ are called stationary points
 - If $f''(a) > 0$ then $x = a$ is a local minimum
 - If $f''(a) < 0$ then $x = a$ is a local maximum

- The derivative of e^x is e^x

- The derivative of $\ln x$ is $\frac{1}{x}$

e.g. If $f(x) = e^x + \ln(2x^3)$ find $f'(x)$

$$f(x) = e^x + \ln 2 + 3\ln x$$

$$f'(x) = e^x + \frac{3}{x}$$

- **Product Rule**

$$\text{If } y = uv \text{ then } \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

- **Quotient Rule**

$$\text{If } y = \frac{u}{v} \text{ then } \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

- **Chain Rule**

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

Find $\frac{dy}{dx}$ given that $\ln(1+x^2)$

Let $u = 1+x^2$ so $y = \ln u$

$$\frac{du}{dx} = 2x \qquad \frac{dy}{du} = \frac{1}{u}$$

$$\frac{dy}{dx} = \frac{2x}{1+x^2}$$

- **Differentiating $\sin x$, $\cos x$ and $\tan x$**

The derivative of **$\sin x$** is **$\cos x$** .

The derivative of **$\cos x$** is **$-\sin x$**

The derivative of **$\tan x$** is **$\sec^2 x$**

- The derivative of $f(ax)$ is $af'(ax)$
- The derivative of $f(ax+b)$ is $af'(ax+b)$

e.g. Find $f'(x)$ given that $f(x) = \sin 3x \cos 2x$

Let $u = \sin 3x$ and $v = \cos 2x$

$$\frac{du}{dx} = 3 \cos 3x \qquad \frac{dv}{dx} = -2 \sin 2x$$

$$\frac{dy}{dx} = (\sin 3x)(-2 \sin 2x) + (\cos 2x)(3 \cos 3x)$$

$$\frac{dy}{dx} = 3 \cos 2x \cos 3x - 2 \sin 2x \sin 3x$$

7 Integration

Key points from C1 and C2

$$\int x^n dx = \frac{x^{n+1}}{n+1} + c$$

$$\int_b^a f(x) dx \quad \text{Gives the area under the graph of } y=f(x) \text{ between } x=a \text{ and } x=b$$

Areas below the x-axis are negative

- Key integrals TO LEARN

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + c \quad \int \frac{1}{x} dx = \ln|x| + c \quad \int \frac{1}{ax+b} dx = \frac{1}{a} \ln|ax+b| + c$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax + c \quad \int \cos ax dx = \frac{1}{a} \sin ax + c$$

- Integration by SUBSTITUTION

e.g. Use the substitution $u = 1-x^2$ to find $\int x\sqrt{1-x^2} dx$

First find du in terms of dx $\frac{du}{dx} = -2x$ so $du = -2x dx$

Rewrite the function in terms of u and du

$$\begin{aligned} \int x\sqrt{1-x^2} dx &= -\frac{1}{2} \int \sqrt{1-x^2} (-2x) dx \\ &= -\frac{1}{2} \int \sqrt{u} du = -\frac{1}{2} \int u^{-\frac{1}{2}} du \end{aligned}$$

Carry out the integration in terms of u

$$= -\frac{1}{2} \left(\frac{2}{3} u^{\frac{3}{2}} \right) + c = -\frac{1}{3} u^{\frac{3}{2}} + c$$

Rewrite the result in terms of x

$$-\frac{1}{3} (1-x^2)^{\frac{3}{2}} + c$$

- Integration by parts

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx + c$$

e.g. Find $\int x e^{5x} dx$

Let $u = x$ and $\frac{dv}{dx} = e^{5x}$

$\frac{du}{dx} = 1$ $v = \frac{1}{5} e^{5x}$

$$\begin{aligned} \int x e^{5x} &= \frac{x}{5} e^{5x} - \int \frac{1}{5} e^{5x} dx \\ &= \frac{x}{5} e^{5x} - \frac{1}{25} e^{5x} + c \end{aligned}$$

- Integrating $\frac{f'(x)}{f(x)}$ (the numerator is a multiple of the derivative of the denominator)

$$\int \frac{f'(x)}{f(x)} = \ln|f(x)| + c$$

e.g. Find $\int \frac{x^3}{x^4 + 1} dx$

The derivative of the denominator, x^4+1 is $4x^3$, so think of the numerator as $\frac{1}{4}(4x^3)$

$$\int \frac{x^3}{x^4 + 1} dx = \frac{1}{4} \int \frac{4x^3}{x^4 + 1} dx = \frac{1}{4} \ln|x^4 + 1| + c$$

$$\left. \begin{aligned} \bullet \int \frac{1}{a^2 + x^2} dx &= \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c \\ \bullet \int \frac{1}{\sqrt{a^2 - x^2}} dx &= \sin^{-1} \left(\frac{x}{a} \right) + c \end{aligned} \right\}$$


STANDARD INTEGRALS TO LEARN

8 Solids of Revolution

- Revolution about the x-axis

The volume of a solid of revolution about the x-axis between $x = a$ and $x = b$ is

given by $\int_a^b \pi y^2 dx$



- Revolution about the y-axis

The volume of a solid of revolution about the y-axis between $y = a$ and $y = b$ is

given by $\int_a^b \pi x^2 dy$


e.g. The region shown is rotated through 2π radians about the y axis.
Find the volume of the solid generated.

First express x^2 in terms of y

$$y = \frac{1}{8}x^3 \Rightarrow 8y = x^3 \Rightarrow 2y^{\frac{1}{3}} = x \Rightarrow x^2 = 4y^{\frac{2}{3}}$$

When $x = 0$ $y = 0$ when $x = 2$ $y = 1$

$$\text{Volume} = \int_0^1 \pi x^2 dy = \int_0^1 4\pi y^{\frac{2}{3}} dy = 4\pi \left[\frac{3y^{\frac{5}{3}}}{5} \right] = \frac{12}{5}\pi$$



9 Numerical Methods

- Change of sign

For an equation $f(x) = 0$, if $f(x_1)$ and $f(x_2)$ have opposite signs and $f(x)$ is continuous between x_1 and x_2 , then a root (solution) of the equation lies between x_1 and x_2

- Staircase and Cobweb Diagrams

If an **iterative** formula (**recurrence relation**) of the form $x_{n+1} = f(x_n)$ converges to a limit, the value of the limit is the x-coordinate of the point of intersection of the graphs $y=f(x)$ and $y = x$

The limit is therefore the solution of the equation $f(x) = x$

A **staircase** or **cobweb** diagram based on the graphs of $y = f(x)$ and $y = x$ illustrates the convergence


e.g. Solve the equation $x^3 - 12x + 12 = 0$

First we will write it in the form $x = f(x)$

$$x^3 + 12 = 12x \Rightarrow \frac{x^3}{12} + 1 = x$$

Plotting the graphs $y = \frac{x^3}{12} + 1$ and $y = x$

the solution is the point of intersection of the two graphs.



We can confirm that there is a point of intersection between $x = 1$ and $x = 2$ by a change of sign the values are substituted.

Substituting $x = 2$ into $y = \frac{x^3}{12} + 1$ gives $y = 1.66\dots$ (shown on the diagram)

Substituting $x = 1.66\dots$ $y = 1.38\dots$

Repeating this the values converge to 1.1157

The solution of $x^3 - 12x + 12 = 0$ is $x = 1.1157$

- The mid-ordinate Rule (Numerical Integration)

Gives an approximation to the area under a graph.


The area is divided into strips of equal width.

The value of the function halfway across each strip (the mid-ordinate) is calculated

Total area = width of strip x sum of mid-ordinates

x	Y
2.5	ln 2.5
3.5	ln 3.5
4.5	ln 4.5
5.5	ln 5.5

$$\begin{aligned} \text{Area} &= 1 \times (\ln 2.5 + \ln 3.5 + \ln 3.5 + \ln 5.5) \\ &= \ln (2.5 \times 3.5 \times 4.5 \times 5.5) \\ &= \ln 216.5625 \\ &= 5.38 \end{aligned}$$



- Simpson's Rule

Gives a more accurate approximation to the area under a graph.

An **even** number of strips of equal width are used.

The ordinates y_0, y_1, y_2, \dots are the values of the function on the vertical edges of the strips. The area is given by

$$\frac{1}{3}h(\text{sum of end ordinates} + 4 \times \text{sum of **odd** ordinates} + 2 \times \text{sum of remaining **even** ordinates})$$